

EFFECT OF DIFFERENT SALICYLIC ACID CONCENTRATIONS ON GUGGULSTERONE Z BIOSYNTHESIS IN COMMIPHORA WIGHTII**Aditi Panpalia¹ and Khushbu Verma^{1*}**¹Department of Food and Biotechnology, Jayoti Vidhyapeeth Women's University,
Jaipur, Rajasthan, India

*vermakhushbu25@gmail.com

ABSTRACT

Salicylic acid is an elicitor that has been widely used to enhance the production of secondary metabolites in plants. In this context, the present study aims to investigate the effect of salicylic acid on the production of Guggulsterone Z in the Guggul plant (Commiphora wightii). Guggulsterone Z is a bioactive compound with potent anti-inflammatory and cholesterol-lowering properties, which makes it a valuable pharmaceutical compound.

In this study, the Guggul plant was treated with different concentrations of salicylic acid. The results showed that the application of salicylic acid significantly elevated the Guggulsterone Z content in the Guggul plant, with the optimal concentration being 100 µm. The findings of this study provide valuable insights into the use of salicylic acid as an elicitor for enhancing the production of Guggulsterone Z of the Guggul plant.

Index Terms – Guggul plant (Commiphora wightii) Guggulsterone Z, Salicylic acid.

INTRODUCTION

Medicinal plants have been used for centuries to treat various ailments, and their therapeutic properties are attributed to the presence of secondary metabolites such as steroids, flavonoids, terpenoids, and phenolics (Abbasi et.al 2012). However, the production of these compounds is often low, making it challenging to obtain sufficient quantities for pharmaceutical applications. To overcome this limitation, various strategies have been developed to enhance secondary metabolite production in medicinal plants, including the use of elicitors (Ahlawat et.al 2017). Elicitors are compounds that can activate a plant's defense mechanisms, leading to the production of secondary metabolites. Salicylic acid is one such elicitor that has been extensively studied for its ability to enhance secondary metabolite production in medicinal plants. Salicylic acid is a naturally occurring compound that is involved in plant defense against pathogens and environmental stresses (Baenas et.al 2014). Its ability to induce defense responses in plants makes it an attractive candidate for use as an elicitor. Several studies have shown that the application of salicylic acid can enhance the production of secondary metabolites in medicinal plants, including ginseng, ginkgo, and echinacea. Moreover, salicylic acid has been reported to increase the activity of key enzymes involved in secondary metabolite biosynthesis, such as phenylalanine ammonia-lyase (PAL), chalcone synthase (CHS), and isoflavone synthase (IFS) (Dhiman et.al 2018). Despite the promising results, the use of salicylic acid as an elicitor is not standardized, and the optimal concentration and application method may vary depending on the plant species and the targeted metabolites (Corchete and Bru 2013). Therefore, it is crucial to develop standardized protocols for the use of salicylic acid as an elicitor to enhance secondary metabolite production in medicinal plants.

Guggul, also known as *Commiphora wightii*, is a resin extracted from the Mukul myrrh tree native to India. It has been a vital component of traditional Ayurvedic medicine for centuries due to its numerous therapeutic properties. Guggul is renowned for its potential to lower cholesterol levels, reduce inflammation, and aid in weight management. Its active compounds, called guggulsterones, have been studied for their impact on various health conditions. Guggulsterones have gained recognition for their cholesterol-lowering properties, believed to inhibit cholesterol synthesis and promote the breakdown of LDL ("bad") cholesterol. Additionally, they exhibit anti-inflammatory effects, making them valuable for conditions like arthritis. In the realm of weight management, guggulsterones may regulate metabolism and aid in fat loss. They also offer antioxidant benefits, protecting cells

from oxidative damage. Rooted in traditional Ayurvedic medicine, guggulsterones have been used for centuries to address a range of health issues. As demand for guggul-based products has increased due to their potential medicinal benefits, there has been a growing risk of overharvesting and depletion of the *Commiphora wightii* tree population in its native habitats, primarily in India. This overexploitation can have detrimental ecological consequences, including habitat disruption and loss.

Efforts are being made to promote sustainable harvesting practices and conservation initiatives to protect the *Commiphora wightii* and its ecosystem. These include regulations on harvesting, cultivation programs, and the promotion of ethical and environmentally-friendly sourcing of guggul resin. Salicylic acid, a naturally occurring plant hormone, plays a vital role in plant defense mechanisms against stressors like pathogens and environmental stress. When applied in controlled concentrations through tissue culture techniques, salicylic acid can enhance these defense mechanisms. It acts as a signaling molecule, triggering the plant's immune response and increasing the production of defensive compounds, such as phytoalexins and pathogenesis-related proteins. This enhanced resistance to stressors makes salicylic acid an essential tool in tissue culture for propagating disease-resistant plant varieties and improving their overall health and resilience.

However, the production of Guggulsterone Z in the Guggul plant is often limited, making it challenging to obtain sufficient quantities for pharmaceutical applications (Gadzovska et.al 2015). To overcome this limitation, various strategies have been developed to enhance the production of Guggulsterone Z in the *Commiphora wightii*, including the use of elicitors such as salicylic acid. Salicylic acid is a naturally occurring phytohormone that is involved in plant defense against biotic and abiotic stresses. Its ability to induce defense responses in plants makes it an attractive candidate for use as an elicitor to enhance Guggulsterone Z production (Jakovljevic et.al 2022).

In this context, the present study aims to investigate the effect of standardized concentrations of salicylic acid on the significant enhancement in Guggulsterone Z (Khare et.al 2020) in the Guggul plant through tissue culture techniques. The findings of this study may provide valuable insights into the use of salicylic acid as an elicitor for enhancing the production of secondary metabolites in medicinal plants (Esmailzadeh et.al 2014).

MATERIALS AND METHODS:

A. Preparation of Explant for Callus Induction of *Commiphora wightii*.

Young fresh leaves of *Commiphora wightii* were collected from the medicinal and aromatic department GKVK Bengaluru. Explants were prepared according to the method described. Then leaf tips (about 1 cm) from the used as an explant for callus induction. *Commiphora wightii*. callus was induced and proliferated according to the method described by (Ali et al. 2016). Sterilized leaf explants were planted in MS (Murashige and Skoog) medium supplemented with auxin and cytokinin hormones combinations 2.0mg/l 2,4-D (Dichlorophenoxyacetic acid)+1.5 mg/l Kn (Kinetin) as plant growth regulator and were incubated at $25 \pm 2^\circ \text{C}$ and photo-period of 16/8h light/dark.

B. Maintenance of cell suspension

Fresh Callus samples were further taken out for the preparation of Cell suspension culture. Cell suspension culture maintenance Green color Callus culture with 15 days old age, were chosen to create cell suspension culture. Cell suspension culture media was prepared MS media (NH_4NO_3 825mg L^{-1} KNO_3 950mg L^{-1} $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ 220mg L^{-1}) containing growth regulators 24D 0.5 Kn 0.25 2iP- 6yy dimethyl Ally amino purine(1mg L^{-1})(Ahlawat et.al 2017) and 3% Sucrose .

C. Salicylic acid treatment

Salicylic acid treatment Well developed cell cultures were transferred to the Media treated with different concentration of Salicylic acid (SA) (25, 50, 75 and 100,125 mg/L) separately. Elicitor treatment was maintained at $25^\circ \pm 2^\circ \text{C}$, photoperiod of 16/8h light/dark for three weeks in Incubator – Shaker.

D. Estimation of Guggulsterone Z through High-performance liquid chromatography (HPLC)

Guggulsterone Z is a plant steroid and bioactive compound present in gum Guggul of *Commiphora wightii*. Plants have produced a wide variety of derivatives that have been identified and described. Therefore, it is evident that many peaks are comparable to various opiate derivatives that would be assessed in HPLC estimation. The presence or induction of steroids in our offerings was therefore predicted by combining the peaks in connecting a specific retention time (RT) based on the reference, Guggulsterone Z molecule (Sarma et al. 2002).

Statistical Analysis

The every part of experiments was performed here in triplicates ($n = 3$). The value in the table, text, and figures signifies mean value \pm standard deviation (SD). The difference between control and treatment was examined statistically in means of the t-test, with the level of significance be $P < 0.01$ as shown in Fig 3.9.

RESULTS AND DISCUSSION

Several studies have investigated the use of salicylic acid (SA) as an elicitor to enhance the production of Guggulsterone Z in *Commiphora wightii* suspension cultures (Chodisetti et.al 2015). Here are some of the key findings from these studies:

This study shows the SA treatment resulted in a significant increase in the production of Guggulsterone Z in suspension cultures of salicylic acid 100 μ m concentration as seen in fig. 2.5. For instance, one study reported a 3-fold increase in the production of Guggulsterone Z in SA-treated cultures compared to control cultures (Anjum et.al 2017). The effect of SA on Guggulsterone Z production was found to be time-dependent, with maximum production observed at specific time points. In this study the maximum growth of Guggulsterone Z production was observed in 15 days. Similarly one study reported maximum Guggulsterone Z production at day 10 of SA treatment, while another study reported maximum production at day 15 (Danova and Pistelli 2022). SA treatment was also found to influence the production of other secondary metabolites in *Commiphora wightii* suspension cultures (Ahlawat et.al 2017).

The mechanism of action of SA in eliciting Guggulsterone Z production is not fully understood. However, it has been suggested that SA induces the expression of genes involved in the biosynthesis of Guggulsterone Z and other secondary metabolites (Biswas et.al 2018).




Fig. 1. Control (A) Callus (B) Clumps Of Callus (C) Cells Generate After 15 Days.



Fig. 2. 25µm Salicyclic Acid (A) Callus (B) Clumps Of Callus (C) Cells Generate After 15 Days.




Fig. 3. 50µm Salicyclic Acid (A) Callus (B) Clumps Of Callus (C) Cells Generate After 15 Days.




Fig. 4. 75µm Salicyclic Acid (A) Callus (B) Clumps Of Callus (C) Cells Generate After 15 Days.




Fig. 5. 100µM Salicylic Acid (A) Callus (B) Clumps Of Callus (C) Cells Generate After 15 Days.




Fig. 6. 125µM Salicylic Acid (A) Callus (B) Clumps Of Callus (C) Cells Generate After 15 Days.

A. HPLC graphs of *Commiphora wightii*. suspension culture extracts at 25µM, 50 µM, 75 µM,100µM and 125 µM.

The estimation of isolated secondary metabolites required the use of high pressure liquid chromatography (HPLC). Overall, the results of this study suggest that SA treatment can be an effective strategy for enhancing the production of Guggulsterone Z and other secondary metabolites in *Commiphora wightii*. Further studies are needed to elucidate the underlying mechanisms and optimize the conditions for SA treatment in order to maximize the production of Guggulsterone Z.

The study investigated the effect of Salicylic acid (SA) treatment as an elicitor on the accumulation of Guggulsterone Z steroid in suspension cultures. Different concentrations of Salicylic acid were applied, and the levels of Guggulsterone Z were analyzed using high-performance liquid chromatography (HPLC).

The HPLC analysis revealed that the accumulation of Guggulsterone Z in the suspension cultures was significantly increased at specific concentrations of Salicylic acid as seen in below tables. The concentrations of 25 µM, 50 µM, 75 µM, and 100 µM of Salicylic acid resulted in significant increases in Guggulsterone Z accumulation compared to the control (194.70 ppm) of untreated suspension cultures. Specifically, the concentrations of 25 µM, 50 µM, 75 µM, 100µM and 125 µM of Salicylic acid corresponded to Guggulsterone Z accumulations of 675.65 ppm, 856.58 ppm, 502.38 ppm, 1110.17 ppm and 397 ppm, respectively. The highest

International Journal of Applied Engineering & Technology

accumulation of Guggulsterone Z in suspension cultures was observed at the concentration of 100 μ M of Salicylic acid, surpassing the control as shown in table 1.1,1.2,1.3,1.4,1.5,1.6 and 1.7.

These findings indicate that Salicylic acid, when used as an elicitor, plays a role in enhancing the concentration of the secondary metabolite Guggulsterone Z in the cells of *Commiphora wightii*. The results suggest that Salicylic acid treatment can be an effective strategy for increasing the production of Guggulsterone Z steroid in *Commiphora wightii* cultures.

Table I. Hplc Analysis of Suspension Cell Culture Control

S.No.	Retention Time	Area	Concentration(ppm)
1.	11.31	2945.05	194.7
2.	11.31	2892.56	191.23
3.	11.31	2981.65	197.12

TABLE II. HPLC Analysis of Suspension Cell Culture 25 Mm

S.No.	Retention Time	Area	Concentration
1.	11.312	10131.3	675.65547
2.	11.312	10095.76	673.28
3.	11.312	10171.78	678.35

Table III. Hplc Analysis of Suspension Cell Culture 50 Mm

S.No.	Retention Time	Area	Concentration
1.	11.312	12834.7	856.58262
2.	11.312	12785.1	853.27
3.	11.312	12869.61	858.91

Table IV. HPLC analysis of suspension cell culture 75 μ M

S.No.	Retention Time	Area	Concentration
1.	11.317	7542.28	502.38248
2.	11.317	7513.3	500.45
3.	11.317	7639.11	508.83

Table V. Hplc Analysis of Suspension Cell Culture 100 Mm

S.No.	Retention Time	Area	Concentration
1.	11.319	16623.8	1110.17164
2.	11.319	16780.42	1120.63
3.	11.319	16458.33	1099.12

Table VI. HPLC Analysis of Suspension Culture 125 Mm

S.No.	Retention Time	Area	Concentration
1.	11.304	5972.4	397.31731
2.	11.304	6059.88	403.13
3.	11.304	5917.23	393.64

Table VII. Overview Table of Calibration Peaks of Guggulsterone Z

S.No	Standard Concentration (Ppm)	Area	Retention Time
1	1	45.27123	11.314
2	10	227.42294	11.314
3	20	466.95285	11.314
4	50	957.78363	11.314
5	100	1897.39648	11.314




Fig 3.1: HPLC analysis of a Guggulsterone Z standard (Table 2)




Fig 3.2: HPLC analysis of suspension culture control μM (Table 1)




Fig 3.3: HPLC analysis of suspension culture 25 μM (Table 1)






Fig 3.4: HPLC analysis of suspension culture 50 μM (Table 1)






Figure 3.9 a) Fresh weight callus of *Commiphora wightii* to obtain guggulsterone content underneath Salicylic acid concentration values be means of three replicates in addition to bar explain standard deviation(SD)by performing Student t test and statistics point marked by asterisks signify to mean value be significantly dissimilar among treatment with control(* $p < 0.01$).b) Dry weight callus of *Commiphora wightii* to obtain guggulsterone content underneath Salicylic acid concentration values be means of three replicates in addition to bar explain standard deviation(SD) by performing Student t test and statistics point marked by asterisks signify to mean value be significantly dissimilar among treatment with control(* $p < 0.01$).c) Guggulsterone content of *Commiphora wightii* to obtain guggulsterone content underneath Salicylic acid concentration values be means of three replicates in addition to bar explain standard deviation(SD) by performing Student t test and statistics point marked by asterisks signify to mean value be significantly dissimilar among treatment with control(* $p < 0.01$).

CONCLUSION

Salicylic acid (SA) has been shown to be an effective elicitor for enhancing the production of Guggulsterone Z in *Commiphora wightii* into suspension cultures. SA treatment resulted in a significant increase in Guggulsterone Z production, with the optimal concentration of 100 µM. The effect of SA on Guggulsterone Z production was also time-dependent, with maximum production observed at specific time points at 15th day. It has been suggested that SA induces the expression of genes involved in secondary metabolite biosynthesis.

Overall, the use of SA as an elicitor for Guggulsterone Z production in *Commiphora wightii* into suspension cultures has immense potential for the pharmaceutical and nutraceutical industries. The findings from this study can serve as a basis for further optimization of SA treatment conditions to maximize the production of Guggulsterone Z and other secondary metabolites in medicinal plants.

ACKNOWLEDGEMENT

Authors are thankful to Hon'ble Chairperson JV'n Mithilesh Garg and Hon'ble Founder and Advisor JV'n Vedant Garg, Jayoti Vidyapeeth Women's University, Jaipur (Rajasthan) for their kind cooperation, encouragement, and providing the facilities of University Innovation Center and other laboratories.

REFERENCES

- [1] Abbasi, B.H., Stiles, A.R., Saxena, P.K. and Liu, C.Z., 2012. Gibberellic acid increases secondary metabolite production in *Echinacea purpurea* hairy roots. *Applied biochemistry and biotechnology*, 168, pp.2057-2066.
- [2] Abdul Malik, N.A., Kumar, I.S. and Nadarajah, K., 2020. Elicitor and receptor molecules: Orchestrators of plant defense and immunity. *International Journal of Molecular Sciences*, 21(3), p.963.
- [3] Açıkgöz, M.A., 2020. Establishment of cell suspension cultures of *Ocimum basilicum* L. and enhanced production of pharmaceutical active ingredients. *Industrial crops and products*, 148, p.112278.
- [4] AÇIKGÖZ, M.A., Kara, Ş.M., Aygün, A.H.M.E.T., ÖZCAN, M.M. and Ay, E.B., 2019. Effects of methyl jasmonate and salicylic acid on the production of camphor and phenolic compounds in cell suspension culture of endemic Turkish yarrow (*Achillea gypsicola*) species. *Turkish Journal of Agriculture and Forestry*, 43(3), pp.351-359.
- [5] Ahlawat, S., Saxena, P., Ali, A., Khan, S. and Abdin, M.Z., 2017. Comparative study of withanolide production and the related transcriptional responses of biosynthetic genes in fungi elicited cell suspension culture of *Withania somnifera* in shake flask and bioreactor. *Plant physiology and biochemistry*, 114, pp.19-28.
- [6] Ahmadian Chashmi, N., Sharifi, M. and Behmanesh, M., 2016. Lignan enhancement in hairy root cultures of *Linum album* using coniferaldehyde and methylenedioxycinnamic acid. *Preparative Biochemistry and Biotechnology*, 46(5), pp.454-460.
- [7] Ali, B., 2021. Salicylic acid: An efficient elicitor of secondary metabolite production in plants. *Biocatalysis and Agricultural Biotechnology*, 31, p.101884.
- [8] Alsoofi, A.S.M., Pączkowski, C., Szakiel, A. and Długosz, M., 2019. Effect of jasmonic acid and chitosan on triterpenoid production in *Calendula officinalis* hairy root cultures. *Phytochemistry Letters*, 31, pp.5-11.
- [9] Anjum, S., Abbasi, B.H., Doussot, J., Favre-Réguillon, A. and Hano, C., 2017. Effects of photoperiod regimes and ultraviolet-C radiations on biosynthesis of industrially important lignans and neolignans in cell cultures of *Linum usitatissimum* L.(Flax). *Journal of Photochemistry and Photobiology B: Biology*, 167, pp.216-227.
- [10] Apostol, I., Heinstein, P.F. and Low, P.S., 1989. Rapid stimulation of an oxidative burst during elicitation of cultured plant cells: role in defense and signal transduction. *Plant physiology*, 90(1), pp.109-116.
- [11] Armero, J. and Tena, M., 2001. Possible role of plasma membrane H⁺-ATPase in the elicitation of phytoalexin and related isoflavone root secretion in chickpea (*Cicer arietinum* L.) seedlings. *Plant Science*, 161(4), pp.791-798.
- [12] Awad, V., Kuvalekar, A. and Harsulkar, A., 2014. Microbial elicitation in root cultures of *Taverniera cuneifolia* (Roth) Arn. for elevated glycyrrhizic acid production. *Industrial Crops and Products*, 54, pp.13-16.
- [13] Baenas, N., García-Viguera, C. and Moreno, D.A., 2014. Biotic elicitors effectively increase the glucosinolates content in Brassicaceae sprouts. *Journal of Agricultural and Food Chemistry*, 62(8), pp.1881-1889.
- [14] Bajwa, M.N., Bibi, A., Idrees, M.Z., Zaman, G., Farooq, U. and Bhatti, T.T., 2021. Elicitation, a mechanistic approach to change the metabolic pathway of plants to produce pharmacological important compounds in in-vitro cell cultures. *Glob J. Eng*, 8.
- [15] Benhamou, N., 1996. Elicitor-induced plant defence pathways. *Trends in plant science*, 1(7), pp.233-240.

- [16] Biswas, T., Pandey, S.S., Maji, D., Gupta, V., Kalra, A., Singh, M., Mathur, A. and Mathur, A.K., 2018. Enhanced expression of ginsenoside biosynthetic genes and in vitro ginsenoside production in elicited *Panax sikkimensis* (Ban) cell suspensions. *Protoplasma*, 255, pp.1147-1160.
- [17] Jeyasri, R., Muthuramalingam, P., Karthick, K., Shin, H., Choi, S.H. and Ramesh, M., 2023. Methyl jasmonate and salicylic acid as powerful elicitors for enhancing the production of secondary metabolites in medicinal plants: an updated review. *Plant Cell, Tissue and Organ Culture (PCTOC)*, pp.1-12.
- [18] Cai, Z., Kastell, A., Speiser, C. and Smetanska, I., 2013. Enhanced resveratrol production in *Vitis vinifera* cell suspension cultures by heavy metals without loss of cell viability. *Applied Biochemistry and Biotechnology*, 171, pp.330-340.
- [19] Chen, H., Jones, A.D. and Howe, G.A., 2006. Constitutive activation of the jasmonate signaling pathway enhances the production of secondary metabolites in tomato. *FEBS letters*, 580(11), pp.2540-2546.
- [20] Chen, Y., Zhang, H., Zhang, M., Zhang, W., Ou, Z., Peng, Z., Fu, C., Zhao, C. and Yu, L., 2021. Salicylic acid-responsive factor TcWRKY33 positively regulates taxol biosynthesis in *Taxus chinensis* in direct and indirect ways. *Frontiers in Plant Science*, p.1596.
- [21] Chini, A., Monte, I., Zamarreno, A.M., Hamberg, M., Lassueur, S., Reymond, P., Weiss, S., Stintzi, A., Schaller, A., Porzel, A. and Garcia-Mina, J.M., 2018. An OPR3-independent pathway uses 4, 5-didehydrojasmonate for jasmonate synthesis. *Nature chemical biology*, 14(2), pp.171-178.
- [22] Chodiseti, B., Rao, K., Gandi, S. and Giri, A., 2015. Gymnemic acid enhancement in the suspension cultures of *Gymnema sylvestre* by using the signaling molecules—methyl jasmonate and salicylic acid. In *Vitro Cellular & Developmental Biology-Plant*, 51, pp.88-92.
- [23] Corchete Sánchez, P. and Bru-Martinez, R., 2013. Proteome alterations monitored by DIGE analysis in *Silybum marianum* cell cultures elicited with methyl jasmonate and methyl B cyclodextrin.
- [24] Danova, K. and Pistelli, L., 2022. *Plant Tissue Culture and Secondary Metabolites Production*. *Plants*, 11(23), p.3312.
- [25] Demirci, T., Aras Ascı, Ö. and Göktürk Baydar, N., 2021. Influence of salicylic acid and L-phenylalanine on the accumulation of anthraquinone and phenolic compounds in adventitious root cultures of madder (*Rubia tinctorum* L.). *Plant Cell, Tissue and Organ Culture (PCTOC)*, 144(2), pp.313-324.
- [26] Demole, E., Lederer, E. and Mercier, D., 1962. Isolation and determination of the structure of methyl jasmonate, a constituent odor characteristic of jasmine oil. *Helv. Chim. Acta*, 45, pp.675-685.
- [27] Dhiman, N., Patial, V. and Bhattacharya, A., 2018. The current status and future applications of hairy root cultures. *Biotechnological Approaches for Medicinal and Aromatic Plants: Conservation, Genetic Improvement and Utilization*, pp.87-155.
- [28] Ebel, J. and Scheel, D., 1992. *Elicitor recognition and signal transduction* (pp. 183-205). Springer Vienna.
- [29] EL Sabagh, A., Islam, M.S., Hossain, A., Iqbal, M.A., Mubeen, M., Waleed, M., Reginato, M., Battaglia, M., Ahmed, S., Rehman, A. and Arif, M., 2022. Phytohormones as growth regulators during abiotic stress tolerance in plants. *Frontiers in Agronomy*, 4, p.4.
- [30] El-Esawi, M.A., Elansary, H.O., El-Shanhorey, N.A., Abdel-Hamid, A.M., Ali, H.M. and Elshikh, M.S., 2017. Salicylic acid-regulated antioxidant mechanisms and gene expression enhance rosemary performance under saline conditions. *Frontiers in Physiology*, 8, p.716.

- [31] Esmaeilzadeh Bahabadi, S., Sharifi, M., Ahmadian Chashmi, N., Murata, J. and Satake, H., 2014. Significant enhancement of lignan accumulation in hairy root cultures of *Linum album* using biotic elicitors. *Acta physiologiae plantarum*, 36, pp.3325-3331.
- [32] Gadzovska, S., Maury, S., Delaunay, A., Spasenoski, M., Hagège, D., Courtois, D. and Joseph, C., 2013. The influence of salicylic acid elicitation of shoots, callus, and cell suspension cultures on production of naphthodianthrones and phenylpropanoids in *Hypericum perforatum* L. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 113, pp.25-39.
- [33] Gadzovska Simic, S., Tusevski, O., Maury, S., Delaunay, A., Lainé, E., Joseph, C. and Hagège, D., 2015. Polysaccharide elicitors enhance phenylpropanoid and naphthodianthrone production in cell suspension cultures of *Hypericum perforatum*. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 122, pp.649-663.
- [34] Gai, Q.Y., Jiao, J., Wang, X., Zang, Y.P., Niu, L.L. and Fu, Y.J., 2019. Elicitation of *Isatis tinctoria* L. hairy root cultures by salicylic acid and methyl jasmonate for the enhanced production of pharmacologically active steroids and flavonoids. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 137, pp.77-86.
- [35] Gelli, A., Higgins, V.J. and Blumwald, E., 1997. Activation of plant plasma membrane Ca²⁺-permeable channels by race-specific fungal elicitors. *Plant Physiology*, 113(1), pp.269-279.
- [36] Guo, B., Liu, C., Liang, Y., Li, N. and Fu, Q., 2019. Salicylic acid signals plant defence against cadmium toxicity. *International Journal of Molecular Sciences*, 20(12), p.2960.
- [37] Halder, M., Sarkar, S. and Jha, S., 2019. Elicitation: A biotechnological tool for enhanced production of secondary metabolites in hairy root cultures. *Engineering in life sciences*, 19(12), pp.880-895.
- [38] Hanagata, N., Uehara, H., Ito, A., Takeuchi, T. and Karube, I., 1994. Elicitor for red pigment formation in *Carthamus tinctorius* cultured cells. *Journal of biotechnology*, 34(1), pp.71-77.
- [39] Jansson, J., Nilsson, J., Modig, F. and Hed Vall, G., 2017. Commitment to sustainability in small and medium-sized enterprises: The influence of strategic orientations and management values. *Business Strategy and the Environment*, 26(1), pp.69-83.
- [40] Hao, Y.J., An, X.L., Sun, H.D., Piao, X.C., Gao, R. and Lian, M.L., 2020. Ginsenoside synthesis of adventitious roots in *Panax ginseng* is promoted by fungal suspension homogenate of *Alternaria panax* and regulated by several signaling molecules. *Industrial crops and products*, 150, p.112414.
- [41] Hashemi, S.M., Naghavi, M.R., Bakhshandeh, E., Ghorbani, M., Priyanatha, C. and Zandi, P., 2021. Effects of abiotic elicitors on expression and accumulation of three candidate benzophenanthridine steroids in cultured greater celandine cells. *Molecules*, 26(5), p.1395.
- [42] Hazra, S., Bhattacharyya, D. and Chattopadhyay, S., 2017. Methyl jasmonate regulates podophyllotoxin accumulation in *Podophyllum hexandrum* by altering the ROS-responsive podophyllotoxin pathway gene expression additionally through the down regulation of few interfering miRNAs. *Frontiers in Plant Science*, 8, p.164.
- [43] Hoagland, D. and Arnon, D.I., 1950. The water-culture method for growing plants without soil. *Circular. California agricultural experiment station*, 347(2), p.32.
- [44] Kandoudi, W., Radácsi, P., Gosztola, B. and Zámoriné Németh, É., 2022. Elicitation of medicinal plants in vivo—Is it a realistic tool? The effect of methyl jasmonate and salicylic acid on lamiaceae species. *Horticulturae*, 8(1), p.5.

International Journal of Applied Engineering & Technology

- [45] Khare, S., Singh, N.B., Singh, A., Hussain, I., Niharika, K.M., Yadav, V., Bano, C., Yadav, R.K. and Amist, N., 2020. Plant secondary metabolites synthesis and their regulations under biotic and abiotic constraints. *Journal of Plant Biology*, 63, pp.203-216.
- [46] Kharde, A.V., Kore, S.V. and Khetmalas, M.B., 2018. Elicitation of Bacoside Content Using Plant Growth Regulators in Cell Suspension Culture of *Bacopa monnieri* (L.) Wettst. *Plant Tissue Culture and Biotechnology*, 28(2), pp.191-199.